

Primary Care Cancer Referral Guidance in Response to COVID-19

Primary Care Cancer Referral Guidance in Response to COVID-19

This guideline has been developed by the SE Primary Care Cancer Reference Group:

Name	Role	Region
Dr Alex Norman	SSCA Primary Care Lead	Surrey and Sussex
Dr Anant Sachdev	TVCA Primary Care Lead	Thames Valley
Dr Tina George	KMCA Primary Care Lead	Kent and Medway
Dr Nigel Watson	WCA Primary Care Lead LMC	Wessex
Dr Chrissie Clayton	Surrey Heartlands Cancer Lead	Surrey
Dr Alex Mancey-Barratt	Sussex Health and Care Cancer Lead	Sussex
Dr Jo Thomson	Coastal West Sussex Cancer Lead	Sussex
Dr Matthew Thomas	East Sussex Cancer Lead	Sussex
Dr Mandeep Ahluwalia	Horsham Mid Sussex Crawley Cancer Lead	Sussex
Dr Richard Brown	Surrey LMC	Surrey and Sussex
Dr Sarnia Ward	Dorset Clinical Lead	Dorset
Dr Tony Newman-Sanders	SE Regional Clinical Advisor for Diagnostics	South East
Keri Torney	SSCA Managing Director	Surrey and Sussex
Stephanie Bell	SSCA Programme Manager	Surrey and Sussex
Becky Clack	Cancer Transformation Manager	Surrey
Jane Cobby	Cancer Transformation Manager	Sussex

1 Purpose of document

The following guidance has been produced to support Primary Care services during the COVID-19 response period. It is intended to guide and support the suspected cancer referral process across the South East Region and to ensure that patients are managed on a case by case basis taking into consideration multiple risk/benefit factors as the pandemic evolves. This document should be viewed as a support for local decision making and should be used alongside National guidance and regionally agreed COVID-19 pathway management guides.

2 Suspected Cancer Referrals 2WW/TWR

Continue to refer suspected cancer as per the normal NG12 referral process but note that trusts have been given latitude in how they are handled. All secondary care providers are being asked to protect patients and staff by reducing the number of visitors to their sites where possible. The South East Cancer Cell is working with clinicians across the region to redesign pathways in response to COVID-19.

Key messages:

- Trusts are encouraged to introduce telephone triage as the first appointment.
- Some patients may have subsequent investigation deferred.
- Full history and pre-referral work up is essential.
- Straight to test may not be an option in the near future on clinical grounds.
- Patients must be made aware of the current NHS pressures and prioritisation and that they may not be seen for some time.

Please Note: Services will be significantly reduced if/when capacity and/or workforce drops to 25% within individual trusts. Any changes to regional guidance will be communicated verbally via the South East Regional Primary Care Reference Group, formally communicated via the ICC and Guidance held on the SE Regional Future NHS COVID-19 platform.

Please Note: in some areas, referrals to multi-diagnostic centres may still be accepted. It is advisable to check with local ICP's for confirmation.

3 Referral responsibilities and management

3.1 Referral management and processing

It is the responsibility of all locality CCGs to ensure that the latest versions of the site specific TWR referral forms reflecting NG12 guidance are available and uploaded on to their respective member GP systems; and old versions removed.

It is the referring GP Practice's responsibility to ensure all TWR referrals are made using the latest version of the cancer site specific referral form reflecting NG12 guidance; and contain all relevant information to enable effective management. This should include the overall COVID-19 risk status for the patient, including frailty score, performance and isolation status for the patient.

It is the responsibility of the referring GP Practice to ensure that the TWR referral form is electronically attached to the e-RS RAS or appointment in a timely way (≤ 24 hours from referral). All patients being referred should have the rationale explained, and the patient should be clear about the importance of them engaging with the pathway.

It is the responsibility of the ICP to ensure that a robust communication function is in place to ensure GP's and Primary care colleagues are kept up to date with local changes during the COVID-19 pandemic. Furthermore, it is advisable that mechanisms such as Advice and Guidance are in place to facilitate communication regarding a patient between secondary and Primary Care.

It is the responsibility of the referring GP Practice to ensure that the patient is aware of the nature of the TWR referral and understands the requirements regarding timeliness and immediate pathway steps:

- Patients should be informed that they may receive a telephone consultation rather than a face to face appointment with the hospital.
- Referrers should inform patients that should they develop any symptoms of COVID-19 including a fever and/or new persistent cough whilst awaiting hospital appointments or tests to contact the hospital and make a different appointment at a later date, after the required period of self-isolation.
- It is the responsibility of the referring GP/clinician to ensure all 'pre-referral' clinical work-up has been completed of relevance to that site specific cancer pathway and in particular that key blood tests and radiological investigations are completed where possible.
- Any patient within an existing pathway wishing to defer appointments, tests or treatment due to COVID-19 related reasons, should be retained by the provider and be visible on the PTL with clear tracking comments; providers must ensure appropriate mechanisms are in place to ensure these patients are not 'lost' in the system and their pathway progressed at a later date. This can be achieved by setting a future tracking date to ensure the patient is reviewed on a tracking work list.

- Patients will NOT be referred back to their GP if unavailable to attend an appointment within the current TWR window.
- Ideally, Trusts should establish site specific RAS on e-RS for all pathways, to enable essential triage processes to be undertaken. This may differ where the clinical complexity of a site specific pathway (diagnostic phase) always dictates a face to face first appointment (consultation); and therefore would not be altered by triage.

4 Safety-netting

- If a patient is unavailable due to COVID-19 isolation reasons (7 days or 14 days – personal symptoms or symptoms within family members), patient will defer accordingly and be picked up within the PTL safety netting process once isolation complete.
- For patients within the vulnerable category (12 weeks but asymptomatic), patients will be encouraged to engage with their pathway teams, and providers should operate care models to reduce hospital contact.

5 Supportive resources for patients with a cancer diagnosis during COVID-19:

1. <https://www.cancercaremap.org/>
2. <https://www.macmillan.org.uk/coronavirus/cancer-and-coronavirus>
3. <https://www.cancerresearchuk.org/about-cancer/cancer-in-general/coronavirus-and-cancer>
4. 'Courageous conversations' training to support staff with difficult conversations. <https://www.macmillan.org.uk/about-us/health-professionals/resources/resources-for-gps.html>

Local services to support cancer patient health and well-being may be available and details should be communicated through ICC teams, CCG's or Health and Wellbeing Boards.

6 Glossary:

CCG	Clinical Commissioning Group
eRS	Electronic Referral System
ICC	Incident Command Centre

ICP	Integrated Care Partnership
TWR	Two Week Referral
PTL	Patient Tracking List
RAS	Referral Assessment Service
NG12	NICE Guideline

